

Amendments to the Claims:

This listing of claims will replace all prior versions, and listings of claims in the application:

Listing of Claims

1. (Currently Amended) A method for the treatment of a renal cell cancer comprising co-administering an anti-tumor antibody directed against the MN antigen and a cytokine to a subject in need thereof, wherein the cytokine is an interferon and is administered continuously or repeatedly in a low-dose form, wherein the low-dose cytokine comprises a dose which is pharmaceutically effective in the absence of NIC CTC toxicity grade 3 or higher.
2. (Currently Amended) A method for the treatment of a renal cell cancer comprising co-administering an anti-tumor antibody directed against the MN antigen and cytokine to a subject in need thereof, wherein the cytokine is an interferon and the method comprises:
 - (a) a first treatment stage comprising administering a low-dose cytokine, and
 - (b) a second treatment stage comprising co-administering the anti-tumor antibody and a low-dose cytokine, and wherein the low-dose cytokine comprises a dose which is pharmaceutically effective in the absence of NIC CTC toxicity grade 3 or higher.
3. (Cancelled)

4. (Previously Presented) The method according to claim 1 comprising a daily administration of a low-dose cytokine.
5. (Cancelled)
6. (Cancelled)
7. (Cancelled)
8. (Currently Amended) The method of claim ~~5~~ 1 wherein the cytokine is IFN- α .
9. (Original) The method of claim 8 wherein the dose of IFN- α is in the range of from 1-10 MIU three times a week.
10. (Currently Amended) The method of claim 1 wherein the cytokine is administered in a ~~substantially~~ constant dose during the treatment.
11. (Previously Presented) The method of claim 1 wherein the cytokine is administered in a variable dose during the treatment.

12. (Previously Presented) The method of claim 1 wherein the cytokine is administered subcutaneously.
13. (Cancelled)
14. (Previously Presented) The method of claim 1 wherein the antitumor antibody is a chimeric or humanized G250 antibody or a fragment thereof.
15. (Previously Presented) The method of claim 1 wherein the antitumor antibody is administered in intervals of from 5-20 days.
16. (Original) The method of claim 2 wherein the first treatment stage comprises 5-20 days.
17. (Original) The method of claim 2 wherein the second treatment stage comprises 50-200 days.